# **Elabscience**®


## Elab Fluor<sup>®</sup> Violet 450 Anti-Human HLA-A,B,C Antibody[W6/32]

### Catalog Number: E-AB-F1130Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Host	Mouse
Isotype	Mouse IgG2a, κ
Clone No.	W6/32
Isotype Control	Elab Fluor <sup>®</sup> Violet 450 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB- F09802Q]
Conjugation	Elab Fluor <sup>®</sup> Violet 450
Conjugation Information	Elab Fluor <sup>®</sup> Violet 450 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 $\mu$ L of antibody per test (million cells in 100 $\mu$ L staining volume or per 100 $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### Data



Human peripheral blood lymphocytes are stained with Elab

Fluor<sup>®</sup> Violet 450 Anti-Human HLA-A,B,C Antibody (Left). Lymphocytes are stained with Elab Fluor<sup>®</sup> Violet 450 Mouse IgG2a,  $\kappa$  Isotype Control (Right).

Preparation & Storage	
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged
	exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	MHC class I;Major Histocompatibility Class I
Uniprot ID	P04439;P01889;P10321
Gene ID	3105

### For Research Use Only

# **Elabscience**®

Background

MHC class I antigens associated with  $\beta$ 2-microglobulin are expressed by all human nucleated cells. MHC class I molecules are involved in presentation of antigens to CD8 + T cells. They play an important role in cell-mediated immune responses and tumor surveillance.