Recombinant Mouse CD159a/KLRC1 Protein (His Tag)

Catalog Number: PKSM040461



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms CD159a;NKG2A;NKG2B

Species Mouse

Expression Host
Sequence
Ala94-Ile244
Accession
AAD24969.1
Calculated Molecular Weight
Observed molecular weight
Tag

HEK293 Cells
Ala94-Ile244
AAD24969.1
19.6 kDa
33-40 kDa
N-His

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

NKG2, also known as NKG2A(CD159A), is a member of the killer cell lectin-like receptor family. This family is a group of transmembrane proteins preferentially expressed in NK cells. Members of this fmaily are characterized by the type II membrane orientation and the presence of a C-type lectin domain. NKG2 contains 1 C-type lectin domain and forms a complex with another family member, KLRD1/CD94. It is expressed only in NK-cells, but not in T-cells or B-cells. It has been shown that NKG2 represents a family of related cDNA clones, designated NKG2A, NKG2B, NKG2C, and NKG2D, which encode type 2 integral membrane proteins (extracellular C-terminus) containing a C-type lectin domain. Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NKG2 functions as a receptor for the

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

Recombinant Mouse CD159a/KLRC1 Protein (His Tag)

Catalog Number:PKSM040461



recognition of MHC class I HLA-E molecules by NK cells and some cytotoxic T-cells.

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com