



A Reliable Research Partner in Life Science and Medicine

Recombinant Histone H3 (Acetyl Lys18) Monoclonal Antibody

catalog number: AN301177L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen A synthetic peptide corresponding to residues around (Lys18) of Human Acety-

Histone H3


HostRabbitIsotypeIgG,κCloneB936PurificationProtein A

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution

WB 1:2000-1:10000

Data



Western Blot with Recombinant Histone H3 (Acetyl Lys18) Monoclonal Antibody at dilution of 1:1000 dilution. Lane A:

C6.

Observed-MW:17 kDa Calculated-MW:15 kDa

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.

For Research Use Only