## Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

# **Mouse CD48 Antibody Pair Set**

Catalog No. E-KAB-0699 Applications ELISA

**Synonyms** CD48;BLAST-1;BCM-1;SLAMF2

## Kit components & Storage

Title	Specifications	Storage
Mouse CD48 Capture Antibody	1 vial, 100 μ g	Store at -20℃. Avoid freeze /
		thaw cycles.
Mouse CD48 Detection Antibody (Biotin)	1 vial, 50 μL	Store at -20℃. Avoid freeze /
		thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items C		Characteristic (E-KAB-0699)	
		Mouse CD48 Capture Antibody	Mouse CD48 Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Mouse CD48 protein	Recombinant Mouse CD48 protein
Information	Swissprot	P18181	
Product details	Reactivity	Mouse	Mouse
	Host	Rabbit	Rabbit
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Affinity purification	Affinity purification
	Specificity	Detects Mouse CD48 in ELISAs.	

For Research Use Only

Tel: 400-999-2100 Web: <a href="www.elabscience.cn">www.elabscience.cn</a> Email: <a href="mailto:techsupport@elabscience.cn">techsupport@elabscience.cn</a>





A Reliable Research Partner in Life Science and Medicine

### **Applications**

Mouse CD48 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4ug/mL	Mouse CD48 Capture	
Capture		Antibody	10 m
ELISA Detection	1:1000-1:10000	Mouse CD48 Detection Antibody (Biotin)	Optical Density
			0.01 100 1000 10000 100000 Mouse CD48 Concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

#### **Background**

Glycosylphosphatidylinositol (GPI)-anchored cell surface glycoprotein that interacts via its N-terminal immunoglobulin domain with cell surface receptors including 2B4/CD244 or CD2 to regulate immune cell function and activation. Participates in T-cell signaling transduction by associating with CD2 and efficiently bringing the Src family protein kinase LCK and LAT to the TCR/CD3 complex. In turn,promotes LCK phosphorylation and subsequent activation. Participates in T-cell signaling transduction by associating with CD2 and efficiently bringing the Src family protein kinase LCK and LAT to the TCR/CD3 complex. In turn,promotes LCK phosphorylation and subsequent activation.

Tel: 400-999-2100 Web: <a href="www.elabscience.cn">www.elabscience.cn</a> Email: <a href="techsupport@elabscience.cn">techsupport@elabscience.cn</a>