

Recombinant Human Interleukin-33/IL-33 Protein (His Tag)

Catalog Number: PKSH032642

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	E.coli-derived Human Interleukin-33; IL-33 protein Ser112-Thr 270, with an C-terminal
	His
Mol_Mass	18.9 kDa
Accession	O95760
Bio-activity	Measure by its ability to binding with recombinant ST2L (IL-33 receptor). The ED ₅₀ for this effect is $<$ 54 ng/mL.Measure by its ability to induce proliferation in D10.G4.1 cells. The ED ₅₀ for this effect is $<$ 0.1 ng/mL. The specific activity of recombinant
	human IL-33 is approximately >4 x10 ⁷ IU/ mg.


Properties	
Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	$< 0.1 \; EU$ per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4.
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

Interleukin-33 (IL-33) belongs to the IL-1 superfamily. IL33 is highly expressed in endothelial venules found in tonsils; Peyer patches and mesenteric lymph nodes; but almost undetectable in placenta. IL33 induces the production of T helper-2 (Th2)-associated cytokines. IL-33 is a cytokine that mediates its biological effects by binding to and signals through IL1RL1/ST2 and IL-1 Receptor Accessory Protein (IL1RAP); activating intracellular molecules in the NF-kB and MAP kinase signaling pathways that drive production of type 2 cytokines from polarized Th2 cells.

For Research Use Only