

Human NCAD Antibody Pair Set

Catalog No. E-KAB-0204 **Applications** ELISA
Synonyms CDH2, CD325, CDHN, CDw325, Cadherin 2 Type 1, N-Cadherin

Kit components & Storage

Title	Specifications	Storage
Human NCAD Capture Antibody	1 vial, 100 µg	Store at -20°C for one year. Avoid freeze / thaw cycles.
Human NCAD Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20°C for one year. Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Product Information

Items		Characteristic (E-KAB-0204)	
		Human NCAD Capture Antibody	Human NCAD Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Human NCAD protein	Recombinant Human NCAD protein
	Swissprot	P19022	
Product details	Reactivity	Human	Human
	Host	Sheep	Sheep
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Human NCAD in ELISAs.	

For Research Use Only

Applications

Human NCAD Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images																
ELISA Capture	0.5-4µg/mL	Human NCAD Capture Antibody	 <p>The graph is a log-log plot of Optical Density versus Human NCAD concentration (ng/mL). The x-axis ranges from 0.01 to 100 ng/mL, and the y-axis ranges from 0.01 to 10. The data points show a clear upward trend, indicating that as the concentration of Human NCAD increases, the optical density also increases. The curve is approximately linear on this log-log scale, suggesting a power-law relationship between the two variables.</p> <table border="1"> <caption>Approximate data points from the standard curve</caption> <thead> <tr> <th>Human NCAD concentration (ng/mL)</th> <th>Optical Density</th> </tr> </thead> <tbody> <tr> <td>0.1</td> <td>0.05</td> </tr> <tr> <td>0.2</td> <td>0.1</td> </tr> <tr> <td>0.5</td> <td>0.2</td> </tr> <tr> <td>1</td> <td>0.4</td> </tr> <tr> <td>2</td> <td>0.8</td> </tr> <tr> <td>5</td> <td>1.5</td> </tr> <tr> <td>10</td> <td>2.5</td> </tr> </tbody> </table>	Human NCAD concentration (ng/mL)	Optical Density	0.1	0.05	0.2	0.1	0.5	0.2	1	0.4	2	0.8	5	1.5	10	2.5
Human NCAD concentration (ng/mL)	Optical Density																		
0.1	0.05																		
0.2	0.1																		
0.5	0.2																		
1	0.4																		
2	0.8																		
5	1.5																		
10	2.5																		
ELISA Detection	1:1000-1:10000	Human NCAD Detection Antibody (Biotin)																	

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Cadherins are a family of transmembrane glycoproteins that mediate calcium-dependent cell-cell adhesion and play an important role in the maintenance of normal tissue architecture. N-cadherin (neural cadherin), also known as CDH2 (cadherin 2), is a classical member of the cadherin superfamily which also includes E-, P-, R-, and B-cadherins. Expression of N-cadherin has been reported on various cell types including neurons, endothelial cells, and cardiac myocytes. N-cadherin has functions in early brain morphogenesis, synaptogenesis, and synaptic plasticity.