A Reliable Research Partner in Life Science and Medicine

## Recombinant Human Transferrin Protein (His Tag)

Catalog Number: PKSH033385

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Species Human

Source HEK293 Cells-derived Human Transferrin protein Val20-Pro698, with an C-terminal His

 Calculated MW
 76.2 kDa

 Observed MW
 85 kDa

 Accession
 AAA61140.1

**Bio-activity** Loaded Recombinant Human Transferrin (PKSH033385) on SA Biosensor, can bind

Recombinant Human TFRC (N-6His) (PKSH033496) with an affinity constant of 99.1

nM as determined in BLI assay.

#### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.


Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.



Loaded Recombinant Human Transferrin (PKSH033385) on SA Biosensor, can bind Recombinant Human TFRC (N-6His) (PKSH033496) with an affinity constant of 99.1 nM as determined in BLI assay.

#### **Background**

# Elabscience®

### Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Serotrans ferrin belongs to transferrin family, and contains 2 transferrin-like domains. The protein is a secreted protein, and expressed by the liver and secreted in plasma. Transferrins are iron binding transport proteins which can bind two Fe3+ ions in association with the binding of an anion. It is responsible for the transport of iron from sites of absorption and heme degradation to those of storage and utilization. Serum transferrin may also have a further role in stimulating cell proliferation.