

APC Anti-Mouse CD64/FcγRI Antibody[X54-5/7.1]

Catalog Number: E-AB-F1186E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	X54-5/7.1
Isotype Control	APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]
Conjugation	APC
Conjugation Information	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
-----	---

Data



C57BL/6 murine bone marrow cells are stained with APC Anti-Mouse CD64 Antibody and FITC Anti-Mouse CD11b Antibody.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	CD64;FcRI;Fcγ1;Fcγr1;IgG Fc receptor I
Uniprot ID	P26151
Gene ID	14129

For Research Use Only

Background

CD64 is a 72 kD single chain type I glycoprotein also known as FcγRI and FcRI. CD64 is a member of the immunoglobulin superfamily and is expressed on monocytes/macrophages, dendritic cells, and mast cells. The expression can be upregulated by IFN-γ stimulation. CD64 binds IgG immune complex. It plays a role in antigen capture, phagocytosis of IgG/antigen complexes, and antibody-dependent cellular cytotoxicity (ADCC).