



A Reliable Research Partner in Life Science and Medicine

Recombinant ZAP-70 Monoclonal Antibody

catalog number: AN300642L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen Recombinant Human ZAP-70 protein

 Host
 Rabbit

 Isotype
 IgG,κ

 Clone
 B577


 Purification
 Protein A

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution

WB 1:2000-1:10000

Data



Western Blot with Recombinant ZAP-70 Monoclonal Antibody at dilution of 1:1000 dilution. Lane A: EL-4-B5 cell lysate.

Observed-MW:70 kDa Calculated-MW:70 kDa

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

This gene encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in this gene cause selective T-cell defect, a severe combined immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been found for this gene.