

Recombinant Human TRIB3/TRB3 Protein (GST Tag)

Catalog Number:PKSH030388



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	C20orf97;NIPK;SINK;SKIP3;TRB3
Species	Human
Expression Host	Baculovirus-Insect Cells
Sequence	Met 1-Gly 358
Accession	NP_066981.2
Calculated Molecular Weight	65.8 kDa
Observed molecular weight	65.8 kDa
Tag	N-GST

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20°C.
Formulation	Supplied as sterile solution of 50mM Tris, 100mM NaCl, pH 8.0, 0.5mM Reduced Glutathione, 10% glycerol, 0.5mM PMSF
Reconstitution	Not Applicable

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Tribbles homolog 3, also known as Neuronal cell death-inducible putative kinase, p65-interacting inhibitor of NF-kappa-B, SINK and TRIB3, is a Nucleus protein which belongs to the protein kinase superfamily and CAMK Ser/Thr protein kinase family and Tribbles subfamily. Highest expression Of TRIB3 is in liver, pancreas, peripheral blood leukocytes and bone marrow. It is also highly expressed in a number of primary lung, colon and breast tumors. TRIB3 is expressed in spleen, thymus, and prostate and is undetectable in other examined tissues, including testis, ovary, small intestine, colon, leukocyte, heart, brain, placenta, lung, skeletal muscle, and kidney. TRIB3 disrupts insulin signaling by binding directly to Akt kinases and blocking their activation. TRIB3 may bind directly to and mask the 'Thr-308' phosphorylation site in AKT1. It binds to ATF4 and inhibits its transcriptional activation activity. TRIB3 interacts with the NF-kappa-B transactivator p65 RELA and inhibits its phosphorylation and thus its transcriptional activation activity. It interacts with MAPK kinases and regulates activation of MAP kinases. It may play a role in programmed neuronal cell death but does not appear to affect non-neuronal cells. TRIB3 does not display kinase activity.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017