

Purified Anti-Mouse CD45.1 Antibody[A20]

catalog number: E-AB-F1184A

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description


Reactivity	Mouse
Immunogen	Recombinant Mouse CD45.1 protein
Host	Mouse
Isotype	Mouse (A.SW) IgG2a, κ
Clone	A20
Purification	>98%, Protein A/G purified
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications

Recommended Dilution

FCM	$\leq 0.2 \mu\text{g}$ per million cells in 100 μL volume
------------	--

Data



HEK293T cells transfected with pcDNA3.1 plasmid encoding Mouse CD45.1 gene were stained with 0.2 μg Purified Anti-Mouse CD45.1 Antibody[A20] (Right) and 0.2 μg Mouse IgG2a, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

CD45 is a 180-240 kD single chain type I membrane glycoprotein also known as leukocyte common antigen (LCA) and T200. It is a tyrosine phosphatase expressed on the plasma membrane of all hematopoietic cells, except erythrocytes and platelets. CD45 is a signaling molecule that regulates a variety of cellular processes including cell growth, differentiation, cell cycle, and oncogenic transformation. CD45 plays a critical role in T and B cell antigen receptor-mediated activation by dephosphorylating substrates including p56Lck, p59Fyn, and other Src family kinases. CD45 non-covalently associates with lymphocyte phosphatase-associated phosphoprotein (LPAP) on T and B lymphocytes. CD45 has been reported to bind galectin-1 and to be associated with several other cell surface antigens including CD1, CD2, CD3, and CD4.

For Research Use Only