

Mergene1000[®] THP-1 Cell-Specific siRNA Transfection Reagent

Cat. No. : 164445

Size: 100 μ L / 0.5mL / 1mL


General Information

Product From	Liquid
Product Color	Colorless
Product Packaging	1 tube
Storage	2-8 $^{\circ}$ C [®]
Expiration Date	18 months
Shipping	Ice bag

Background

Mergene1000[®] THP-1 Cell-Specific siRNA Transfection Reagent is a new and stable siRNA transfection reagent with efficient RNA compression ability. It can quickly and efficiently transfect RNA into cells, while forming a stable complex to effectively protect RNA from nuclease degradation. It is specifically formulated for use with THP-1 cells, achieving high transfection efficiency. The reagent is distinguished by its low toxicity, excellent stability, ease of operation, and high reproducibility.

Product Operation Flowchart



Usage Steps

To transfect THP-1 cells, follow the steps outlined below. Using 24-well plates as an example, mix Mergene1000[®] THP-1 Cell-Specific siRNA Transfection Reagent (μ L) with siRNA (pmol) at a ratio of 1:20. This ratio can be adjusted between 1:15 and 1:30 according to the situation. For other sizes of culture plates or dishes, refer to the recommended transfection amounts provided in Table 1.

1. Cell seeding

Before transfection, ensure that the cells are healthy and cell viability exceeding 90%. Add 500 μ L RPMI-1640 (PM150110) + 10% FBS + 0.05 mM β -mercaptoethanol + 1% P/S (PB180120) medium to each well, and start transfection when the final cell density reaches $2-4 \times 10^5$ cells/mL/well.

2. Preparation of the transfection complex

- (1) Prepare two sterile centrifuge tubes. In one tube, add 20 pmol siRNA and RPMI-1640 medium (PM150110) to a final volume of 10 μ L, and gently mix by pipetting. In the other tube, add 1.0 μ L of Mergene1000® THP-1 Cell-Specific siRNA Transfection Reagent and 9.0 μ L of RPMI-1640 medium (PM150110) to a final volume of 10 μ L, blow and mix.

Note: The above is the amount of preparation for each well of cells. Please calculate the required volumes based on your specific experimental conditions and requirements.

- (2) Mix the above two equal volume diluents, gently mix by pipetting, and incubate for 20 minutes at room temperature.

3. Cell transfection

- (1) Add the prepared 20 μ L transfection complex dropwise to the cells and mixed, incubated at 37°C with 5% CO₂ for culture.
- (2) After 18-48 hours of incubation, detect gene expression.

Table 1. Reference dosage of THP-1 cells transfection in different culture vessel

Culture Vessel	Area	Cell Seeding Density	Inoculation Medium	Diluted Final Volume	siRNA Transfection	
					Reagent Amount	siRNA Amount
96-well	0.3 cm ²	1-4 $\times 10^4$ cells/well	200 μ L	2 \times 5 μ L	0.2 μ L	4 pmol
24-well	2.0 cm ²	2-4 $\times 10^5$ cells/well	500 μ L	2 \times 10 μ L	1.0 μ L	20 pmol
12-well	4.0 cm ²	4-8 $\times 10^5$ cells/well	1 mL	2 \times 20 μ L	2.0 μ L	40 pmol
6-well	10.0 cm ²	1-2 $\times 10^6$ cells/well	2 mL	2 \times 50 μ L	4.0 μ L	80 pmol
6 cm	20.0 cm ²	2-4 $\times 10^6$ cells/well	5 mL	2 \times 0.1 mL	8.0 μ L	160 pmol
10 cm	60.0 cm ²	0.6-1.2 $\times 10^7$ cells/well	15 mL	2 \times 0.3 mL	24.0 μ L	480 pmol

Note: The usage amounts provided in the table are for reference only. The exact amount of siRNA used with Mergene1000® THP-1 Cell-Specific siRNA Transfection Reagent should be optimized according to the cell conditions and other experimental parameters.

Notes

1. This product is only used for scientific research or further research, not for diagnosis and treatment
2. The cell inoculation amount and transfection ratio provided above are based on experiments conducted with THP-1 cells and are for reference only. The specific experimental dosage should be adjusted according to the actual conditions.
3. The product is transported with ice bag and stored at 2-8°C. It can be aliquoted and stored upon use to avoid multiple prolonged openings of the lid.
4. RPMI-1640 medium should be prepared separately for the dilution of siRNA and transfection reagents.
5. After transfection, there is no need to remove the transfection complex or replace with fresh culture medium. The actual operation can be based on the cell status, after transfection culture 4-6 hours to choose to change the medium.
6. The use of high purity siRNA is helpful to obtain higher transfection efficiency. Before transfection, ensure that siRNA gene silencing expression does not affect cell viability.

7. The experimental process utilized RNA-free and pyrogen-free materials, such as centrifuge tubes, pipette tips, and buffers.
8. For in vitro transfection only.
9. For your safety and health, please wear experimental clothes and wear disposable gloves aseptic operation.

Experimental Results Show (For reference only)




Figure 1. Bright-field and fluorescence images of THP-1 cells transfected with FAM-siRNA using Mergene1000[®] THP-1 Cell-Specific siRNA Transfection Reagent.




Figure 2. THP-1 cells were transfected with NC-siRNA and GAPDH-siRNA using Mergene1000[®] THP-1 cell-specific siRNA transfection reagent, and the relative GAPDH expression was detected by qPCR.