

## APC Anti-Human Myeloperoxidase Antibody[1B10]

Catalog Number: AN00925E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description


Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	1B10
Isotype Control	APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]
Conjugation	APC
Conjugation Information	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications

### Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
-----	---

### Data



Intracellular staining of normal human peripheral blood cells with APC Anti-Human Myeloperoxidase Antibody[1B10] (filled gray histogram) or APC Mouse IgG1, κ Isotype Control (empty black histogram). Cells in the granulocytes gate were used for analysis.

### Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

### Antigen Information

Alternate Names	MPO
Uniprot ID	P05164
Gene ID	4353

### For Research Use Only

## Background

Myeloperoxidase (MPO) is a heterotetrameric protein consisting of two 60 kD heavy units and two 12 kD light units. A lysosomal enzyme, MPO is able to catalyze the production of hypochlorous acid, a potent microbicidal agent, from hydrogen peroxide and chloride anion during the neutrophil respiratory burst. MPO is a major enzyme involved in the inflammatory responses of polymorphonuclear leucocytes. MPO is localized to the azurophilic granules of mature granulocytes and monocytes and is also expressed in some acute myeloid leukemia cells.