

Recombinant CD68 Monoclonal Antibody

catalog number: **AN300590P**

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description


Reactivity	Rat
Immunogen	Recombinant Rat CD68 Protein
Host	Rabbit
Isotype	IgG
Clone	B525
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS

Applications


Recommended Dilution

IHC-P	1:1000-1:4000
--------------	---------------

Data



Immunohistochemistry of paraffin-embedded rat lung using CD68 Monoclonal Antibody at dilution of 1:2000.



Immunohistochemistry of paraffin-embedded rat liver using CD68 Monoclonal Antibody at dilution of 1:2000.

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

Background

Macrosialin, also known as CD68 and Gp11, is a single-pass type I membrane protein which belongs to the LAMP family. CD68 is highly expressed by blood monocytes and tissue macrophages. It is also expressed in lymphocytes, fibroblasts and endothelial cells. CD68 is expressed in many tumor cell lines which could allow them to attach to selectins on vascular endothelium, facilitating their dissemination to secondary sites. CD68 plays a role in phagocytic activities of tissue macrophages, both in intracellular lysosomal metabolism and extracellular cell-cell and cell-pathogen interactions. It is a commonly used marker for macrophages. However, a number of studies have shown that CD68 antibodies react with other hematopoietic and non-hematopoietic cell types, suggesting that CD68 may not be a macrophage-specific antigen. CD68 binds to tissue- and organ-specific lectins or selectins, allowing homing of macrophage subsets to particular sites. Rapid recirculation of CD68 from endosomes and lysosomes to the plasma membrane may allow macrophages to crawl over selectin-bearing substrates or other cells.

For Research Use Only