


Recombinant Cyclin B1 Monoclonal Antibody

catalog number: **AN301834L**

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description	
Reactivity	Human
Immunogen	Recombinant human Cyclin B1 fragment
Host	Rabbit
Isotype	Rabbit IgG
Clone	C371456
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.
Applications	Recommended Dilution
WB	1:500-1:1000
IF	1:50
IP	1:50-1:100

Data




Western Blot with Cyclin B1 Monoclonal Antibody at dilution of 1:1000. Lane 1: Jurkat, Lane 2: HeLa, Lane 3: HT-29

Observed-MW:55 kDa
Calculated-MW:55 kDa



Immunoprecipitation analysis using anti-Cyclin B1 Monoclonal Antibody. Western blot was performed from the immunoprecipitate using Cyclin B1 Monoclonal Antibody at a dilution of 1:100. Lane 1: 5% Input, Lane 2: Cyclin B1 Monoclonal Antibody, Lane 3: Rabbit monoclonal IgG Isotype

Observed-MW:55 kDa
Calculated-MW:55 kDa



Immunofluorescent analysis of (100% Ice-cold methanol) fixed Jurkat cells using anti-Cyclin B1 Monoclonal Antibody at dilution of 1:50.

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.3

Shipping

Ice bag

Background

Cyclins are a family of proteins that activate specific cyclin-dependent kinases required for progression through the cell cycle. The entry of all eukaryotic cells into mitosis is regulated by activation of cdc2/cdk1 at the G2/M transition. This activation is a multi-step process that begins with the binding of the regulatory subunit, cyclin B1, to cdc2/cdk1 to form the mitosis-promoting factor (MPF). MPF remains in the inactive state until phosphorylation of cdc2/cdk1 at Thr161 by cdk activating kinase (CAK) and dephosphorylation of cdc2/cdk1 at Thr14/Tyr15 by cdc25C. Five cyclin B1 phosphorylation sites (Ser116, 126, 128, 133, and 147) are located in the cytoplasmic retention signal (CRS) domain and are thought to regulate the translocation of cyclin B1 to the nucleus at the G2/M checkpoint, promoting nuclear accumulation and initiation of mitosis.

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.3