

IL-18/IGIF, Mouse, Recombinant

Cat. No. : GPCK311

产品信息

物种	Mouse
表达宿主	E.coli
序列信息	MNFGRLHCTTAVIRNINDQVLFVDKRQPVFEDMTDIDQSASEPQTRLIIYMYKDSEVRGLAVTL SVKDSKMSTLSCKNKIISFEEMDPPENIDDIQSDLIFFQKRVPGHNMKEFESSLYEGHFLACQK EDDAFKLILKKKDENGDKSVMFTLTNLHQS with polyhistidine tag at the C-terminus.
检索号	P70380.2
标签	His-tag at the C-terminus
分子量	19.1 kDa
有效期	12 months
生物活性	Measure by its ability to induce IFN gamma secretion in KG-1 cells. The ED50 for this effect is < 0.5 µg/mL.

产品特性

内毒素 (EU/µg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1 × PBS, pH 8.0.
复溶	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Interleukin-18 (IL-18) is a cytokine that belongs to the IL-1 superfamily and is produced by macrophages and other cells. IL-18 works by binding to the interleukin-18 receptor, and together with IL-12 it induces cell-mediated immunity following infection with microbial products like lipopolysaccharide (LPS). After stimulation with IL-18, natural killer (NK) cells and certain T cells release another important cytokine called interferon-γ (IFN-γ) or type II interferon that plays an important role in activating the macrophages or other cells. The combination of this cytokine and IL12 has been shown to inhibit IL-4 dependent IgE and IgG1 production, and enhance IgG2a production in B cells. IL-18 binding protein (IL18BP) can specifically interact with this cytokine, and thus negatively regulate its biological activity.

