

Recombinant PTEN Monoclonal Antibody

catalog number: **AN301026L**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;Mouse;Rat
Immunogen	Recombinant Human PTEN protein
Host	Rabbit
Isotype	IgG, κ
Clone	5A1
Purification	Protein A
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution


WB	1:1000-1:5000
IHC	1:200-1:1000

Data




Western Blot with Recombinant PTEN Monoclonal Antibody at dilution of 1:1000 dilution. Lane A: MCF7 cells.

Observed-MW:56 kDa
Calculated-MW:47 kDa



Immunohistochemistry of paraffin-embedded human pancreas tissue using Recombinant PTEN Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded rat pancreas tissue using Recombinant PTEN Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

For Research Use Only

Phosphatase and tensin homolog (PTEN) Homo sapiens This gene was identified as a tumor suppressor that is mutated in a large number of cancers at high frequency. The protein encoded by this gene is a phosphatidylinositol-3, 4,5-trisphosphate 3-phosphatase. It contains a tensin like domain as well as a catalytic domain similar to that of the dual specificity protein tyrosine phosphatases. Unlike most of the protein tyrosine phosphatases, this protein preferentially dephosphorylates phosphoinositide substrates. It negatively regulates intracellular levels of phosphatidylinositol-3,4,5-trisphosphate in cells and functions as a tumor suppressor by negatively regulating AKT/PKB signaling pathway. The use of a non-canonical (CUG) upstream initiation site produces a longer isoform that initiates translation with a leucine, and is thought to be preferentially associated with the mitochondrial inner membrane.