

CCL2/MCP-1, Swine, Recombinant**Cat. No. : GPCK351****产品信息**

| | |
|------|---|
| 物种 | Swine |
| 表达宿主 | E.coli |
| 序列信息 | QPDAINSPVTCCYTLTSKKISMQRRLMSYRRVTSSKCPKEAVIFKTIAGKEICAEPKQKWVQDSI SHLDKKNQTPKP with polyhistidine tag at the N-terminus. |
| 检索号 | P42831.1 |
| 标签 | His-tag at the N-terminus |
| 分子量 | 9.42 kDa |
| 有效期 | 12 months |

产品特性

| | |
|-------------|--|
| 内毒素 (EU/μg) | < 0.1 |
| 保存 | Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months. |
| 运输 | Ambient temperature or ice pack. |
| 制剂 | The protein was lyophilized from a 0.2 μm filtered solution containing 1 × PBS, pH 7.4. |
| 复溶 | It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein. |

背景介绍

CCL2, also known as MCP-1, is belonging to the CC ? chemokine family. CCL2 can be identified in endothelial cells, smooth muscle cells and monocytes as the results of reaction to several atherogenic stimulants, such as CD40 ligand, (IL-1β) and oxidized low density lipoprotein, interleukin-1β platelet derived growth factor (PDGF). Recent study shows that in vivo MCP1 have several critical roles in atherosclerosis. Additionally, MCP-1 has been proved involving in monocytic infiltration of tissues during several inflammatory diseases, and has been implicated in macrophage-mediated tumor growth inhibition in mice. In addition, CCL2 has been shown to have direct effects on tumor cells in an autocrine and paracrine fashion in multiple cancers, including sarcoma, lung, cervix, ovary, breast, and prostate.

