



A Reliable Research Partner in Life Science and Medicine

Purified Anti-Human HLA-DQ Antibody[1A3]

catalog number: AN004210P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human HLA-DQ protein

Host Mouse

Isotype Mouse IgG2a, κ

Clone 1A3

Purification >98%, Protein A/G purified


Buffer Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer.

Dialyze to completely remove the stabilizer prior to labeling.

Applications Recommended Dilution

FCM $2 \mu g/mL(0.5 \times 10^6 - 1 \times 10^6 \text{ cells})$

Data



Human peripheral blood lymphocytes were stained with 0.2 μ g Purified Anti-Human HLA-DQ Antibody[1a3] (Right) and 0.2 μ g Mouse IgG2a, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD19 PE-conjugated Monoclonal Antibody.

Preparation & Storage

Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

Background

HLA-DQ is heterodimeric cell surface glycoprotein comprised of a 27 kD α (heavy) chain and a 32 kD β (light) chain. In contrast to other MHC class II molecules, both chains of HLA-DQ are polymorphic and the α chain shows a high degree of polymorphism. It is expressed on B cells, activated T cells, monocytes/macrophages, dendritic cells, and other non-professional APCs. In hematopoietic development, HLA-DR and DP are expressed first, followed by HLA-DQ. Variations in the HLA gene expression are crucial to graft survival.

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.0