Elabscience®

ANAPC2 Polyclonal Antibody


catalog number: E-AB-92831

Note: Centrifuge before opening to ensure complete recovery of vial contents.

1:50-1:200


Description	
Reactivity	Mouse;Rat
Immunogen	A synthetic peptide of human ANAPC2
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000

IHC Data




Western blot analysis of extracts of various cell lines using ANAPC2 Polyclonal Antibody at 1:1000 dilution.

Observed-MV:102 kDa Calculated-MV:93 kDa



Immunohistochemistry of paraffin-embedded rat lung using ANAPC2 Polyclonal Antibody at dilution of 1:300 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded rat spleen using ANAPC2 Polyclonal Antibody at dilution of 1:300 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining

protocol.

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the
	temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Elabscience®

A large protein complex, termed the anaphase-promoting complex (APC), or the cyclosome, promotes metaphaseanaphase transition by ubiquitinating its specific substrates such as mitotic cyclins and anaphase inhibitor, which are subsequently degraded by the 26S proteasome. Biochemical studies have shown that the vertebrate APC contains eight subunits. The composition of the APC is highly conserved in organisms from yeast to humans. The product of this gene is a component of the complex and shares sequence similarity with a recently identified family of proteins called cullins, which may also be involved in ubiquitin-mediated degradation.

For Research Use Only