

Elab Fluor® 647 Anti-Human CD305 Antibody[NKTA255]


Catalog Number: GFH00330M

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1
Clone No.	NKTA255
Isotype Control	Elab Fluor® 647 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792M]
Conjugation	Elab Fluor® 647
Conjugation Information	Elab Fluor® 647 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide.

Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (millie cells in 100 µL staining volume or per 117 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of normal human peripheral blood cells with FITC

Anti-Human CD3 Antibody and Elab Fluor® 647 Anti-Human CD305 Antibody[NKTA255] (left) or Elab Fluor® 647 Mouse IgG1, κ Isotype Control (right). Cells in the lymphocytes gate were used for analysis.

Preparation & Storage	
Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	LAIR-1
Uniprot ID	Q6GTX8
Gene ID	3903

For Research Use Only

Tel: 400-999-2100

Web: www.elabscience.cn

Email: techsupport@elabscience.cn

Rev. V1.5

Background

CD305, also known as LAIR-1 (leukocyte-associated Ig-like receptor-1), is a 40 kD type I transmembrane glycoprotein. LAIR-1 and LAIR-2 are the only members of the LAIR family in the Ig superfamily. LAIR-1 is an inhibitory molecule characterized by ITIMs in the cytoplasmic domain and collagen ligands. CD305 is expressed on NK cells, T cells, B cells, monocytes, dendritic cells, eosinophils, basophils and mast cells. LAIR-1 functions in the inhibition of cell cytotoxicity, cell activation, proliferation and differentiation.