
## Recombinant Human VEGF121 protein(His Tag)

## Catalog Number: PDMH100373

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| SpeciesHumanSourceHEK293 Cells-derived Human VEGF121 protein Met1-Arg147, with an C-terminalCalculated MW17.2 kDaObserved MW14 kDa;18 kDaAccessionP15692-9Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU/mg of the protein as determined by the LAL method | al His   |
|--|----------|
| Calculated MW17.2 kDaObserved MW14 kDa; 18 kDaAccessionP15692-9Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Findotoxin< 1.0 EU/mg of the protein as determined by the LAL method   | al His   |
| Observed MW14 kDa;18 kDaAccessionP15692-9Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU/mg of the protein as determined by the LAL method  |          |
| AccessionP15692-9Bio-activityNot validated for activityPropertiesPurity>95% as determined by reducing SDS-PAGE.Endotoxin<1.0 EU/mg of the protein as determined by the LAL method  |          |
| Bio-activity Not validated for activity   Properties Purity > 95% as determined by reducing SDS-PAGE.   Fndotoxin < 1.0 EU/mg of the protein as determined by the LAL method   |          |
| Properties   Purity > 95% as determined by reducing SDS-PAGE.   Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method   Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots            |          |
| Purity> 95% as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU/mg of the protein as determined by the LAL method   |          |
| Endotoxin< 1.0 EU/mg of the protein as determined by the LAL methodStorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20<br>°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots  |          |
| StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots   |          |
| °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots   |          |
|  | 0 to -80 |
|  | of       |
| reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.  |          |
| Shipping This product is provided as lyophilized powder which is shipped with ice packs  | 5.       |
| <b>Formulation</b> Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%   |          |
| Mannitol.  |          |
| <b>Reconstitution</b> It is recommended that sterile water be added to the vial to prepare a stock solu  | tion of  |
| 0.5 mg/mL. Concentration is measured by UV-Vis.  |          |

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. Induces endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces permeabilization of blood vessels. Binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. NRP1/Neuropilin-1 binds isoforms VEGF-165 and VEGF-145. Isoform', 'binds to KDR but does not activate downstream signaling pathways, does not activate angiogenesis and inhibits tumor growth. Binding to NRP1 receptor initiates a signaling pathway needed for motor neuron axon guidance and cell body migration, including for the caudal migration of facial motor neurons from rhombomere 4 to rhombomere 6 during embryonic development. Also binds the DEAR/FBXW7-AS1 receptor.