

Recombinant Human TGFB2 Protein

Catalog Number: PKSH033139

Note: Centrifuge before opening to ensure complete recovery of vial contents.



Description

Species	Human
Source	HEK293 Cells-derived Human TGFB2 protein Ala303-Ser414
Calculated MW	12.7 kDa
Observed MW	12 kDa
Accession	P61812
Bio-activity	Measured by its ability to inhibit the IL-4-dependent proliferation of TF-1 cells. The ED ₅₀ for this effect is 30-180pg/ml.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.01 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 4mM HCl. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



Measured by its ability to inhibit the IL-4-dependent proliferation of TF-1 cells. The ED₅₀ for this effect is 30-180pg/ml.

Background

Transforming growth factor beta-2 (TGF-β2) is a secreted protein which belongs to the TGF-beta family. It is known as a cytokine that performs many cellular functions and has a vital role during embryonic development. The precursor is cleaved into mature TGF-beta-2 and LAP, which remains non-covalently linked to mature TGF-beta-2 rendering it inactive. It is an extracellular glycosylated protein. It is known to suppress the effects of interleukin dependent T-cell tumors. Defects in TGFB2 may be a cause of non-syndromic aortic disease (NSAD).

For Research Use Only