SMARCA4 Polyclonal Antibody

catalog number: E-AB-70117



Note: Centrifuge before opening to ensure complete recovery of vial contents.

110	C C	TO THE	\mathbf{n}	\mathbf{n}
De		ш	JUL	υш

Reactivity Human; Mouse; Rat

Immunogen KLH conjugated Synthetic peptide corresponding to Mouse BRG1

Host Rabbit Isotype IgG

Purification Affinity purification
Conjugation Unconjugated


buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 1% protein

protectant and 50% glycerol.

Applications Recommended Dilution


WB 1:500-1:2000 **IHC** 1:200-1:600

Data



Western Blot analysis of various samples using SMARCA4 Polyclonal Antibody at dilution of 1:750.


Observed-MV:185 kDa Calculated-MV:185 kDa



Immunohistochemistry analysis of paraffin-embedded mouse testis using SMARCA4 Polyclonal Antibody at dilution of 1:300.



Immunohistochemistry analysis of paraffin-embedded Human tonsil using SMARCA4 Polyclonal Antibody at dilution of 1:300.



Immunohistochemistry analysis of paraffin-embedded rat kidney using SMARCA4 Polyclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

SMARCA4 Polyclonal Antibody

catalog number: E-AB-70117



The protein encoded by this gene is a member of the SWI/SNF family of proteins and is similar to the brahma protein of Drosophila. Members of this family have helicase and ATPase activities and are thought to regulate transcription of certain genes by altering the chromatin structure around those genes. The encoded protein is part of the large ATP-dependent chromatin remodeling complex SNF/SWI, which is required for transcriptional activation of genes normally repressed by chromatin. In addition, this protein can bind BRCA1, as well as regulate the expression of the tumorigenic protein CD44. Mutations in this gene cause rhabdoid tumor predisposition syndrome type 2. Multiple transcript variants encoding different isoforms have been found for this gene.