

Recombinant Human TSPAN1 Protein (aa 110-211, Fc Tag)

Catalog Number: PKSH030743

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human TSPAN1 protein Tyr110-Asn211, with an N-terminal rFc
Calculated MW	39.1 kDa
Observed MW	43-53 kDa
Accession	O60635
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

TSPAN1 belongs to the transmembrane 4 superfamily, also known as the tetraspanin family. Tetraspanins have four hydrophobic domains, intracellular N- and C-termini and two extracellular domains. Tetraspanins act as scaffolding proteins, anchoring multiple proteins to one area of the cell membrane. They also mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. TSPAN1 interacts with human thiamine transporter-1 (hTHTR-1). HTHTR-1 contributes to intestinal thiamine uptake, and its function is regulated at both the transcriptional and posttranscriptional levels. TSPAN1 and hTHTR-1 colocalize in human intestinal epithelial HuTu-80 cells. Coexpression of TSPAN1 in these cells led to a significant decrease in the rate of degradation of hTHTR-1 compared with cells expressing the hTHTR-1 alone; in fact the half-life of the TSPAN1 protein was twice longer in the former cell type compared with the latter cell type.