

PDGF-BB/PDGF2, Human, Recombinant**Cat. No. : GPCK067****产品信息**

物种	Human
表达宿主	E.coli
序列信息	Ser82-Thr190
检索号	P01127
分子量	12.42 kDa
有效期	12 months
生物活性	Measured in a cell proliferation assay using BALB/c 3T3 cells. The ED50 for this effect is 5-20 ng/mL.

产品特性

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 μm filtered solution of 20 mM NaAC-HAC, 4% Mannitol, pH 4.5.
复溶	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

Platelet-Derived Growth Factor Subunit B (PDGFB) belongs to the PDGF/VEGF Growth Factor family. Platelet-Derived Growth Factor is a potent mitogen for cells of mesenchymal origin. PDGFB can exist either as a homodimer (PDGF-BB) or as a heterodimer with the platelet-Derived Growth Factor alpha polypeptide (PDGF-AB), where the dimers are connected by disulfide bonds. Mutations in this gene are associated with meningioma. Binding of PDGFB to its Receptor elicits a variety of cellular responses. In addition, PDGFB is released by platelets upon wounding and plays an important role in stimulating adjacent cells to grow and thereby heals the wound.

