Elabscience®

LIMK1 Polyclonal Antibody

catalog number: E-AB-93157

Note: Centrifuge before opening to ensure complete recovery of vial contents.

1:500-1:2000

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human LIMK1
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution

WB

Data



Western blot analysis of extracts of various cell lines using

LIMK1 Polyclonal Antibody at 1:1000 dilution.

Observed-MW:73 kDa

Calculated-MW:33 kDa/68 kDa/70 kDa/72 kDa

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the
	temperature recommended.

Background

There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. LIMK1 is a serine/threonine kinase that regulates actin polymerization via phosphorylation and inactivation of the actin binding factor cofilin. This protein is ubiquitously expressed during development and plays a role in many cellular processes associated with cytoskeletal structure. This protein also stimulates axon growth and may play a role in brain development. LIMK1 hemizygosity is implicated in the impaired visuospatial constructive cognition of Williams syndrome. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

For Research Use Only