



A Reliable Research Partner in Life Science and Medicine

Mouse MMP-8 Antibody Pair Set

Catalog No. E-KAB-0121 Applications ELISA

Synonyms MMP8, CLG1, HNC, PMNL-CL, Neutrophil Collagenase

Kit components & Storage

Title	Specifications	Storage
Mouse MMP-8 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Mouse MMP-8 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0121)	
		Mouse MMP-8 Capture Antibody	Mouse MMP-8 Detection Antibody (Biotin)
Immunogen	Immunogen	Recombinant Mouse MMP-8 protein	Recombinant Mouse MMP-8 protein
Information	Swissprot	O70138	
Product details	Reactivity	Mouse	Mouse
	Host	Rabbit	Rabbit
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Protein A & Antigen Affinity	Protein A & Antigen Affinity
	Specificity	Detects Mouse MMP-8 in ELISAs.	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com



A Reliable Research Partner in Life Science and Medicine

Applications

Mouse MMP-8 Sandwich ELISA Assav:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Mouse MMP-8 Capture Antibody	
Capture			Ais Ais
ELISA Detection	1:1000-1:10000	Mouse MMP-8 Detection Antibody (Biotin)	0.00 0.01 0.00 100 100 100 100 100 100 1
			Mouse MMP-8 concentration(ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. However, the enzyme encoded by this gene is stored in secondary granules within neutrophils and is activated by autolytic cleavage. Its function is degradation of type I, II and III collagens. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com