

# Recombinant Human 4-1BB/CD137 protein (His tag)

Catalog Number:PDMH100148



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description

|                                    |  |
|------------------------------------|--|
| <b>Synonyms</b>                    | CD137;ILA;TNFRSF9;4-1BB ligand receptor;CDw137;T-cell antigen 4-1BB homolog;T-cell antigen ILA |
| <b>Species</b>                     | Human  |
| <b>Expression Host</b>             | HEK293 Cells   |
| <b>Sequence</b>                    | Met1-Gln186  |
| <b>Accession</b>                   | Q07011   |
| <b>Calculated Molecular Weight</b> | 20.4 kDa   |
| <b>Observed molecular weight</b>   | 30 kDa   |
| <b>Tag</b>                         | C-His  |

## Properties

|                       |   |
|-----------------------|---|
| <b>Purity</b>         | > 95 % as determined by reducing SDS-PAGE.  |
| <b>Endotoxin</b>      | Please contact us for more information.   |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.   |
| <b>Formulation</b>    | Lyophilized from sterile PBS, pH 7.4.<br>Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual.           |
| <b>Reconstitution</b> | Please refer to the printed manual for detailed information.  |

## Background

Tumor necrosis factor receptor superfamily member 9(TNFRSF9); also known as CD137 and 4-1BB; is an inducible T cell surface protein belonging to the tumor necrosis factor receptor superfamily. It is a single-pass type I membrane protein which contains 4 TNFR-Cys repeats. The human and mouse proteins share 60% amino acid sequence identity. CD137 is expressed by mesenchymal cells; including endothelial cells; chondrocytes; and cells of the central nervous system. CD137 is also broadly expressed by cells of the human immune system; is broadly expressed by cells of the human immune system; including activated CD8+ and CD4+ T cells; activated natural killer (NK) cells; follicular dendritic cells (FDCs) and monocytes. CD137 has diverse roles in the immune response; the one key function is to promote the survival of both T cells and dendritic cells by binding the cognate ligand CD137L (4-1BBL).

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017