

Osteocalcin Monoclonal Antibody(Capture)

catalog number: AN001510P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Rat
Immunogen	Recombinant Rat Osteocalcin protein expressed by E.coli
Host	Mouse
Isotype	Mouse IgG1
Clone	10A5
Purification	Protein A/G Purification
Conjugation	Unconjugated
buffer	Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

Applications

Recommended Dilution

ELISA Capture	2-8 µg/mL
----------------------	-----------

Data



Sandwich ELISA-Recombinant Rat Osteocalcin protein standard curve. Background subtracted standard curve using Osteocalcin antibody(AN001510P)(Capture), Osteocalcin antibody(AN001500P)(Detector) in sandwich ELISA. The reference range value for Recombinant Rat Osteocalcin protein is 0.78-50 ng/mL.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

Osteocalcin Monoclonal Antibody(Capture)

catalog number: AN001510P



The carboxylated form is one of the main organic components of the bone matrix, which constitutes 1-2% of the total bone protein: it acts as a negative regulator of bone formation and is required to limit bone formation without impairing bone resorption or mineralization. The carboxylated form binds strongly to apatite and calcium.

The uncarboxylated form acts as a hormone secreted by osteoblasts, which regulates different cellular processes, such as energy metabolism, male fertility and brain development. Regulates energy metabolism by acting as a hormone favoring pancreatic beta-cell proliferation, insulin secretion and sensitivity and energy expenditure. Uncarboxylated osteocalcin hormone also promotes testosterone production in the testes: acts as a ligand for G protein-coupled receptor GPRC6A at the surface of Leydig cells, initiating a signaling response that promotes the expression of enzymes required for testosterone synthesis in a CREB-dependent manner. Also acts as a regulator of brain development: osteocalcin hormone crosses the blood-brain barrier and acts as a ligand for GPR158 on neurons, initiating a signaling response that prevents neuronal apoptosis in the hippocampus, favors the synthesis of all monoamine neurotransmitters and inhibits that of gamma-aminobutyric acid (GABA). Osteocalcin also crosses the placenta during pregnancy and maternal osteocalcin is required for fetal brain development.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine
Tel: 400-999-2100

Email: techsupport@elabscience.cn

Web: www.elabscience.cn

Rev. V1.5