Purified Anti-Human CD4 Antibody[13B8.2]

catalog number: AN004390P



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human CD4 protein

HostRabbitIsotypeRabbit IgG, κ Clone13B8.2

Purification >98%, Protein A/G purified


Conjugation Unconjugated

buffer PBS, pH 7.2. Contains 0.05% proclin 300.

Applications Recommended Dilution

FCM $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$

Data



Human peripheral blood lymphocytes were stained with 0.2 μg Purified Anti-Human CD4 Antibody[13B8.2] (Right) and 0.2 μg Rabbit IgG1, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Rabbit IgG Secondary Antibody.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Order now, ship in 3 days

Background

CD4 is an approximately 55 kDa type I membrane glycoprotein that is expressed predominantly on most thymocytes and a subset of mature T lymphocytes. In humans, CD4 is also expressed to a lesser extent on monocytes and macrophage related cells. Human CD4 cDNA encodes a 458 amino acid (aa) residue precursor protein with a 25 aa residue signal peptide, a 371 aa residue extracellular region containing four immunoglobulin homology domains, a 24 aa residue transmembrane domain and a 38 aa residue cytoplasmic domain.

For Research Use Only