

# Recombinant Human CDK2 Protein (E.coli, His Tag)

Catalog Number:PDEH100009



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	Cyclin-Dependent Kinase 2;Cell Division Protein Kinase 2;p33 Protein Kinase;CDK2;CDKN2
<b>Species</b>	Human
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Met 1-Leu298
<b>Accession</b>	P24941
<b>Calculated Molecular Weight</b>	36.1 kDa
<b>Observed molecular weight</b>	34 kDa
<b>Tag</b>	N-His

## Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4., 5% trehalose, 5% mannitol, 0.01% tween-80. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the print
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 90 % as determined by reducing SDS-PAGE.

## Background

Cyclin-dependent kinase 2 (CDK2) belongs to the cyclin-dependent kinase of Ser/Thr protein kinase. CDK2 acts as a catalytic subunit of the cyclin dependent kinase complex, whose activity is restricted to the G1-S phase of the cell cycle, it is essential for the G1/S transition. The kinase activity of CDK2 can be regulated by the association with a cyclin subunit, its phosphorylation state and CDK inhibitors. The activation of the CDK2/cyclin complex requires the phosphorylation of Thr160 and the dephosphorylation of Try14 and Tyr15. The inhibition of CDK2-cyclin complex can also be attributed to association with p27Kip1 and p21Waf1/Cip1. The activation of CDK2 has been shown to be necessary for apoptosis of quiescent cells, such as neurons, thymocytes and endothelial cells.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017