

## Elab Fluor® 700 Anti-Mouse CD172a/SIRPα Antibody[P84]

Catalog Number: E-AB-F1286M1

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Mouse
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG1, κ
<b>Clone No.</b>	P84
<b>Isotype Control</b>	Elab Fluor® 700 Rat IgG1, κ Isotype Control[HRPN] [Product E-AB-F09822M1]
<b>Conjugation</b>	Elab Fluor® 700
<b>Conjugation Information</b>	Elab Fluor® 700 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 719 nm (e.g., a 725/40 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications

### Recommended usage

<b>FCM</b>	Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

### Data



Staining of C57BL/6 murine bone marrow cells with FITC

Anti-Mouse/Human CD11b Antibody[M1/70] and Elab Fluor® 700 Anti-Mouse CD172a/SIRPα Antibody[P84](left) or Elab

Fluor® 700 Rat IgG1, κ Isotype Control(right). Total viable cells were used for analysis.

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	BIT;CD172 antigen-like family member A;P84;PTPNS1;SHPS-1;SIRPα
<b>Uniprot ID</b>	Q64314
<b>Gene ID</b>	19261

### For Research Use Only

## Background

CD172a, also known as SIRP $\alpha$ , is a type I transmembrane protein with one V-set Ig-like and two C-set Ig-like domains in the extracellular portion, and two ITIM motifs and a proline-rich region in the cytoplasmic tail. CD172a is expressed by monocytes, macrophages, myeloid cells, and neuronal tissue. The phosphorylation of SIRP $\alpha$  ITIMs induces the recruitment and activation of the tyrosine phosphatases PTPN6 and PTPN11, resulting in the negative regulation of several biological processes. The ligands of CD172a are CD47, SP-A, and SP-D.