
Elabscience®

Elab Fluor[®] Violet 450 Anti-Human CD5 Antibody[HISM2]

Catalog Number: E-AB-F1313Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Reactivity	Human	
Host	Mouse	
Isotype	Mouse IgG1, ĸ	
Clone No.	HISM2	
Isotype Control	Elab Fluor [®] Violet 450 Mouse IgG1, к Isotype Control[MOPC-21] [Product E-AB- F09792Q]	
Conjugation	Elab Fluor [®] Violet 450	
Conjugation Information	Elab Fluor [®] Violet 450 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).	
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.	
Applications	Recommended usage	
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.	



Human peripheral blood lymphocytes are stained with APC

Anti-Human CD19 Antibody and Elab Fluor[®] Violet 450 Anti-Human CD5 Antibody[HISM2] (Left). Lymphocytes are stained with APC Anti-Human CD19 Antibody and Elab Fluor

[®] Violet 450 Mouse IgG1, к Isotype Control (Right).

Preparation & Storage		
Storage	Keep as concentrated solution.	
	This product can be stored at 2-8°C for 12 months. Please protec exposure to light and do not freeze.	ted from prolonged
Shipping	Ice bag	
Antigen Information		
Alternate Names	CD5;LEU1;Lymphocyte antigen T1/Leu-1;T-cell surface glycoprotein CD5	
Uniprot ID	P06127	
For Research Use Only	/	
Toll-free: 1-888-852-8623	Tel: 1-832-243-6086	Fax: 1-832-243-6017
Web:www.elabscience.com	Email:techsupport@elabscience.com	Rev. V1.6

Elabscience®

Gene ID Background 921

CD5 is a 67 kD single chain type I glycoprotein also known as Leu-1, Ly-1 and T1. It is a member of the scavenger receptor superfamily found on T cells, thymocytes, B cell subsets, chronic B lymphocytic leukemia (B-Cells), and peripheral blood dendritic cell s. CD5 modulates T and B cell receptor signaling, thymocyte maturation, and T-B cell interactions upon binding to ligands such as CD72.