

A Reliable Research Partner in Life Science and Medicine

Recombinant COPS3 Monoclonal Antibody

catalog number: AN301497L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human;Rat;Mouse

Immunogen Recombinant human COPS3 fragment


HostRabbitIsotypeIgG, κCloneA192


Purification Protein Apurified

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution WB 1:500-1:1000 IHC 1:50-1:100 IF 1:50


Data






Western Blot with COPS3 Monoclonal Antibody at dilution of Immunohistochemistry of paraffin-embedded Human colon 1:1000. Lane 1: HeLa, Lane 2: 293, Lane 3: N2a using COPS3 Monoclonal Antibody at dilution of 1:100.


Observed-MW:44 kDa Calculated-MW:44 kDa





Immunohistochemistry of paraffin-embedded Mouse cardiac muscle using COPS3 Monoclonal Antibody at dilution of 1:100.

Immunohistochemistry of paraffin-embedded Rat cardiac muscle using COPS3 Monoclonal Antibody at dilution of 1:100.



Immunofluorescent analysis of (100% Ice-cold methanol) fixed HeLa cells using anti-COPS3 Monoclonal Antibody at dilution of 1:50.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com



Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

Component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (UbI) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the UbI ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, lkappaBalpha/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the UbI system, respectively.

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.0