A Reliable Research Partner in Life Science and Medicine

Recombinant Mouse ECM1 Protein (His Tag)

Catalog Number: PKSM040729

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Mouse

Source HEK293 Cells-derived Mouse ECM1 protein Met 1-Glu 559, with an C-terminal His

 Mol_Mass
 62.5 kDa

 Accession
 Q61508-1

Bio-activity Measured by its ability to bind human MMP-9 in a functional ELISA.

Properties

Purity > 88 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU per } \mu\text{g}$ of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 88 % as determined by reducing SDS-PAGE.

Background

Extracellular matrix protein 1 (ECM1) is a secreted glycoprotein and playing a pivotal role in endochondral bone formation, angiogenesis, and tumour biology. Three splice variants have been identified: ECM1a (540 aa) is most widely expressed, with highest expression in the placenta and heart; ECM1b (415 aa) is differentiation-dependent expressed and found only in tonsil and associated with suprabasal keratinocytes; ECM1c (559 aa) accounts for approximately 15% of skin ECM1. Although ECM1 is not tumor specific, is significantly elevated in many malignant epithelial tumors and is suggested as a possible trigger for angiogenesis, tumor progression and malignancies. It also has been shown to regulate endochondral bone formation, skeletal development and tissue remodeling.

For Research Use Only

Fax: 1-832-243-6017