

NADPH oxidase 4 (NOX4) Polyclonal Antibody

catalog number: **E-AB-93356**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Mouse;Rat
Immunogen	Recombinant fusion protein of human NADPH oxidase 4
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

WB	1:500-1:2000
IHC	1:50-1:100


Data




Western blot analysis of extracts of various cell lines using NADPH oxidase 4 Polyclonal Antibody at 1:1000 dilution.

Observed-MW:67 kDa

Calculated-MW:6 kDa/25-31 kDa/58-66 kDa




Immunohistochemistry of paraffin-embedded rat lung using NADPH oxidase 4 Polyclonal Antibody at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Western blot analysis of extracts of various cell lines using NADPH oxidase 4 Polyclonal Antibody at 1:1000 dilution.

Observed-MW:67 kDa

Calculated-MW:6 kDa/25-31 kDa/58-66 kDa



Immunohistochemistry of paraffin-embedded rat kidney using NADPH oxidase 4 Polyclonal Antibody at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

This gene encodes a member of the NOX-family of enzymes that functions as the catalytic subunit the NADPH oxidase complex. The encoded protein is localized to non-phagocytic cells where it acts as an oxygen sensor and catalyzes the reduction of molecular oxygen to various reactive oxygen species (ROS). The ROS generated by this protein have been implicated in numerous biological functions including signal transduction, cell differentiation and tumor cell growth. A pseudogene has been identified on the other arm of chromosome 11. Alternative splicing results in multiple transcript variants.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.7