EGF Polyclonal Antibody(Capture/Detector)

catalog number: AN000770P



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| - | | | | | | ٠, | | |
|--------------|---|---|----|----|----|----|----|---|
| D | Α | | C | PТ | m | ŤΤ | U. | m |
| \mathbf{L} | v | 0 | v. | ш | עו | UΙ | U. | |

Reactivity Mouse

Immunogen Recombinant Mouse EGF protein expressed by Mammalian

Host Rabbit
Isotype Rabbit IgG

Purification Antigen Affinity Purification


Conjugation Unconjugated

buffer Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

Applications Recommended Dilution

ELISA Capture 2-8 μg/mL **ELISA Detector** 0.1-0.4 μg/mL

Data



Sandwich ELISA-Recombinant Mouse EGF protein standard curve. Background subtracted standard curve using EGF antibody(AN000770P)(Capture), EGF antibody(AN000770P) (Detector) in sandwich ELISA. The reference range value for Recombinant Mouse EGF protein is 15.63-1000 pg/mL.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

This gene encodes a member of the epidermal growth factor superfamily. The encoded preproprotein is proteolytically processed to generate the 53-amino acid epidermal growth factor peptide. This protein acts a potent mitogenic factor that plays an important role in the growth, proliferation and differentiation of numerous cell types. This protein acts by binding with high affinity to the cell surface receptor, epidermal growth factor receptor. Defects in this gene are the cause of hypomagnesemia type 4. Dysregulation of this gene has been associated with the growth and progression of certain cancers. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed.

For Research Use Only