# **Elabscience**®

## **GSTA3** Polyclonal Antibody


### catalog number: E-AB-15087

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Human
Immunogen	Recombinant protein of human GSTA3
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:25-1:100

#### Data





Western Blot analysis of Human testis tissue using GSTA3 Polyclonal Antibody at dilution of 1:250 Immunohistochemistry of paraffin-embedded Human brain using GSTA3 Polyclonal Antibody at dilution of 1:30



Immunohistochemistry of paraffin-embedded Human prostate

cancer using GSTA3 Polyclonal Antibody at dilution of 1:30

#### Preparation & Storage

Storage Shipping Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

### For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w .elabscience.com</u>

# **Elabscience**®

Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two distinct supergene families. These enzymes are involved in cellular defense against toxic, carcinogenic, and pharmacologically active electrophilic compounds.At present, eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta.This gene encodes a glutathione S-transferase belonging to the alpha class genes that are located in a cluster mapped to chromosome 6.Genes of the alpha class are highly related and encode enzymes with glutathione peroxidase activity.However, during evolution, this alpha class gene diverged accumulating mutations in the active site that resulted in differences in substrate specificity and catalytic activity.

For Research Use Only