Recombinant Mouse Cd80 Protein(FC Tag)

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Catalog Number: PDMM100052



Description Species Mouse Source Mammalian-derived Mouse CD80/B7-1 proteins Val38-Asn246, with an C-terminal Fc Mol Mass 47.9 kDa O00609-1 Accession **Bio-activity** Not validated for activity **Properties** >90% as determined by reducing SDS-PAGE. Purity Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. Shipping This product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Formulation Mannitol. Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data

kDa	М	R
80 60	11	8
40	-	
30	-	
20	-	
12	٣	

SDS-PAGE analysis of Mouse CD80/B7-1 proteins, 2 µg/lane of Recombinant Mouse CD80/B7-1 proteins was resolved with an SDS-PAGE under reducing conditions, showing bands at 47.9 KD

Background

The B-lymphocyte activation antigen B7-1 (referred to as B7), also known as CD80, is a member of cell surface immunoglobulin superfamily and is expressed on the surface of antigen-presenting cells including activated B cells, macrophages and dendritic cells. As costimulatory ligands, B7-1 which exists predominantly as dimer and the related protein B7-2, interact with the costimulatory receptors CD28 and cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) expressed on T cells, and thus constitute one of the dominant pathways that regulate T cell activation and tolerance, cytokine production, and the generation of CTL. The B7/CD28/CTLA4 pathway has the ability to both positively and negatively regulate immune responses. CD80 is thus regarded as promising therapeutic targets for autoimmune diseases and various carcinomas.

For Research Use Only