

Recombinant Human CD47 Protein (Fc Tag)

Catalog Number:PKSH030857



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	IAP;MER6;OA3
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Pro139
Accession	NP_942088.1
Calculated Molecular Weight	40.7 kDa
Observed molecular weight	48-60 kDa
Tag	C-hFc
Bioactivity	1. Immobilized human SIRPA-His at 10 µg/ml (100 µl/well) can bind human CD47-Fc, The EC50 of human CD47-Fc is 10.1-23.5 ng/ml. 2. Immobilized mouse SIRPA-His at 10 µg/ml (100 µl/well) can bind human CD47-Fc, The EC50 of human CD47-Fc is 0.05-0.13 µg/ml. 3. Immobilized human SIRPG-His at 10 µg/ml (100 µl/well) can bind human CD47-Fc, The EC50 of human CD47-Fc is 0.58-1.34 µg/ml.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CD47 contains 1 Ig-like V-type (immunoglobulin-like) domain and is a receptor for the C-terminal cell binding domain of thrombospondin. It may play a role in membrane transport and signal transduction. CD47 is also a membrane protein;

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human CD47 Protein (Fc Tag)

Catalog Number:PKSH030857



which is involved in the increase in intracellular calcium concentration that occurs upon cell adhesion to extracellular matrix. It is very broadly distributed on normal adult tissues; as well as ovarian tumors; being especially abundant in some epithelia and the brain. CD47 may play a role in membrane transport and/or integrin dependent signal transduction. It may prevent premature elimination of red blood cells. It also may be involved in membrane permeability changes induced following virus infection. By acting as an adhesion receptor for THBS1 on platelets; CD47 plays a role in both cell adhesion and in the modulation of integrins. It also plays an important role in memory formation and synaptic plasticity in the hippocampus.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017