

## McCoy's 5A (with HEPES) Complete Medium (20)

Cat. No. : PM150711C

Size : 125mL×4

### General Information

<b>Product Form</b>	Liquid
<b>Concentration</b>	Ready-to-use
<b>Components</b>	McCoy's 5A, with HEPES[PM150711]+20% Nutrient+1% Supplement
<b>Bacterial detection</b>	Negative
<b>Fungal detection</b>	Negative
<b>Mycoplasmal detection</b>	Negative
<b>Endotoxin level</b>	< 3 EU/mL
<b>Shipping</b>	Ice bag
<b>Storage</b>	2-8°C, Shading Light
<b>Expiry date</b>	6 months

### Product Introduction

McCoy's 5A medium was initially used specifically for the culture of Novikoff Hepatoma cells. Unlike other medium, McCoy's 5A medium contains reduced glutathione, bacterial peptone and high concentrations of glucose. McCoy's 5A is widely used in the culture of primary cells, such as bone marrow, skin, gingiva, kidney, spleen, lung, rat embryo, omentum and so on. In addition, McCoy's 5A medium is also used for tissue biopsy culture, the establishment of cell line, and the culture of some lymphocytes and more difficult cells.

### Guidelines for use

1. Procell's cell culture media undergoes strict quality control to ensure sterility, but may get contaminated during use. Follow these guidelines for sterile handling to avoid contamination.
2. Always wipe your gloved hands and work area with 70% ethanol.
3. Wipe the outside of the containers, flasks, plates, and dishes with 70% ethanol before placing them in the cell culture hood.
4. Use sterile pipette tips and pipettes to work with liquids, and use each pipette tip only once to avoid cross contamination. Do not unwrap sterile pipettes until they are ready to be used. Keep pipettes and tips within the clean work area.
5. Do not talk while performing sterile procedures and perform your cell culture as efficiently and carefully as possible to minimize contamination.

### Quality control

Standard evaluations for cell culture media are pH, osmolality, endotoxins and sterility testing for liquid products, cell growth experiments.

### Notes

1. This product is only used for scientific research or further research, not for diagnosis and treatment.
2. This product is sterilized by 0.1 µm filtration.
3. It is necessary to pay attention to the aseptic operation and avoid the contamination during the culture.
4. It is not suitable for long time storage at room temperature.
5. This product is a ready-to-use medium. If there is no special need, don't add serum, penicillin and streptomycin. It can be used directly.

Procell system®

Procell system®

Procell system®

Procell system®