A Reliable Research Partner in Life Science and Medicine

Recombinant Human OX-2/MOX1/CD200 Protein (His Tag)

Catalog Number: PKSH032840

Note: Centrifuge before opening to ensure complete recovery of vial contents.

\mathbf{r}							
H)	es	C	m	n	т	ſΠ	ï

Species Human

Source HEK293 Cells-derived Human OX-2; MOX1; CD200 protein Gln31-Gly232, with an C-

terminal His

Calculated MW 23.5 kDa
Observed MW 35-50 kDa
Accession P41217

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.Formulation Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CD200 is a transmembrane immunoregulatory protein that belongs to the immunoglobulin superfamily. It contains one Ig like V type domain and one Ig like C2 type domain in its extracelluar domain. CD200 is widely but not ubiquitously expressed. Its receptor (CD200R) is restricted primarily to mast cells; basophils; macrophages; and dendritic cells; which suggests myeloid cell regulation as the major function of CD200. CD200 and CD200R associate via their respective N-terminal Ig-like domains. In myeloid cells; CD200R initiates inhibitory signals following receptor-ligand contact. In T cell s; CD200 functions as a co-stimulatory molecule independent of the CD28 pathway. In addition; CD200 also plays an important role in prevention of graft rejection; autoimmune diseases and spontaneous abortion.

For Research Use Only