Recombinant Human Apolipoprotein C-II/APOC2 Protein (His Tag)


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Catalog Number: PKSH032084



Description **Species** Human 10.0 kDa Mol Mass Accession AAP35354.1 Not validated for activity **Bio-activity Properties** > 95 % as determined by reducing SDS-PAGE. Purity Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method. Store at $< -20^{\circ}$ C, stable for 6 months. Please minimize freeze-thaw cycles. Storage This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel Shipping packs. Upon receipt, store it immediately at $< -20^{\circ}$ C. Formulation Supplied as a 0.2 µm filtered solution of PBS, 50% Glycerol, pH 7.4. Reconstitution Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

APOC2 activates the lipoprotein lipase in capillaries, which hydrolyzes triglycerides and thus provides free fatty acids for cells. APOC2 is component of the very low density lipoprotein (VLDL) fraction in plasma. It is also an activator of several triacylglycerol lipases. The association of APOC2 with plasma chylomicrons, VLDL, and HDL is reversible, a function of the secretion and catabolism of triglyceride-rich lipoproteins, and changes rapidly. Defects in APOC2 are the cause of hyperlipoproteinemia type 1B (HLPP1B) which characterized by hypertriglyceridemia, xanthomas, and increased risk of pancreatitis and early atherosclerosis.

For Research Use Only