

## Recombinant Human TNF- $\alpha$ protein (His Tag)

Catalog Number: PDMH100384

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

Species	Human
Source	HEK293 Cells-derived Human TNF- $\alpha$ protein Val77-Leu233, with an C-terminal His
Calculated MW	17.2 kDa
Observed MW	15 kDa
Accession	P01375
Bio-activity	Not validated for activity

### Properties

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Tumor Necrosis Factor- $\alpha$  (TNF- $\alpha$ ) is secreted by macrophages, monocytes, neutrophils, T-cells, and NK-cells following stimulation by bacterial LPS. Cells expressing CD4 secrete TNF- $\alpha$  while cells that express CD8 secrete little or no TNF- $\alpha$ . Synthesis of TNF- $\alpha$  can be induced by many different stimuli including interferons, IL2, and GM-CSF. The clinical use of the potent anti-tumor activity of TNF- $\alpha$  has been limited by the proinflammatory side effects such as fever, dose-limiting hypotension, hepatotoxicity, intravascular thrombosis, and hemorrhage. Designing clinically applicable TNF- $\alpha$  mutants with low systemic toxicity has been of intense pharmacological interest. Human TNF- $\alpha$  that binds to murine TNF-R55 but not murine TNF-R7, exhibits retained anti-tumor activity and reduced systemic toxicity in mice compared with murine TNF- $\alpha$ , which binds to both murine TNF receptors. Based on these results, many TNF- $\alpha$  mutants that selectively bind to TNF-R55 have been designed. These mutants displayed cytotoxic activities on tumor cell lines in vitro and have exhibited lower systemic toxicity in vivo. Recombinant Human TNF- $\alpha$  High Active Mutant differs from the wild-type by amino acid substitution of amino acids 1-7 with Arg8, Lys9, Arg10 and Phe157. This mutant form has been shown to have increased activity with less inflammatory side effects in vivo.