

## Recombinant Human PECAM1/CD31 Protein (His Tag)

Catalog Number: PDMH100398

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	Mammalian-derived Human PECAM1 protein Met1-Lys601, with an C-terminal His
<b>Calculated MW</b>	66.0 kDa
<b>Observed MW</b>	100 kDa
<b>Accession</b>	P16284
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 90% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



SDS-PAGE analysis of Human PECAM1/CD31 proteins, 2 µg/lane of Recombinant Human PECAM1/CD31 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 100 kDa.

### Background

#### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

CD31, also known as platelet endothelial cell adhesion molecule-1 (PECAM-1), is a 130 kDa heavily glycosylated transmembrane protein belonging to the immunoglobulin (Ig) superfamily of cell adhesion molecules. CD31 is highly expressed on endothelial cells and at a lower level on platelets, granulocytes, macrophages, dendritic cells, T and B cells, and natural killer (NK) cells. It is involved in cell adhesion and is required for transepithelial migration of leukocytes (TEM). CD31 is composed of an extracellular domain (ECD) of 574 amino acids containing six Ig-like domains, a transmembrane domain, and a 118 aa cytoplasmic domain. The latter undergoes alternative splicing which generates multiple isoforms showing altered adhesive properties compared to full length CD31. The human CD31 ECD shares 63% and 61% aa sequence identity with mouse and rat CD31, respectively. CD31 acts as a homophilic receptor through its extracellular domain and is involved in downstream signaling via its cytoplasmic domain. This domain contains highly conserved ITIM motifs which, once tyrosine phosphorylated, recruit and activate the signaling molecules Src and SHP-2. The resulting inhibition of TCR signaling increases the activation threshold of T cells, thus reinforcing peripheral tolerance and preventing development of autoimmunity. CD31 additionally regulates immune responses by acting as a key inhibitory receptor in dendritic cell development. Besides its role in TEM, CD31 appears to regulate T cell trafficking through a complex coordination of endothelial cell junctions and T cell extravasation.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017