A Reliable Research Partner in Life Science and Medicine

# Recombinant Mouse PDGFRa/CD140a Protein (Fc Tag)

Catalog Number: PKSM041120

Note: Centrifuge before opening to ensure complete recovery of vial contents.

## Description

Species Mouse

Source HEK293 Cells-derived Mouse PDGFRa/CD140a protein Leu25-Glu524, with an C-

terminal Fc

Calculated MW 83.2 kDa
Observed MW 120-150 kDa
Accession P26618

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### **Background**

Platelet-derived growth factor receptors (PDGFR) are cell surface tyrosine kinase receptors for members of the platelet-derived growth factor (PDGF) family. The PDGF family consists of PDGF-A, -B, -C and -D, which form either homo- or heterodimers (PDGF-AA, -AB, -BB, -CC, -DD). The four PDGFs are inactive in their monomeric forms. PDGFs bind to the protein tyrosine kinase receptors PDGF receptor- $\alpha$  and - $\beta$ . These two receptor isoforms dimerize upon binding the PDGF dimer, leading to three possible receptor combinations, namely - $\alpha\alpha$ , - $\beta\beta$  and - $\alpha\beta$ . PDGFR $\alpha$  and PDGFR $\beta$  are members of the class III RTK family. Inappropriate PDGFR $\alpha$  and PDGFR $\beta$  signaling has been linked to a number of proliferative disorders.

### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017