


## Recombinant MMP-2 Monoclonal Antibody

catalog number: AN300112P

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

| Description          |                                                     |
|----------------------|-----------------------------------------------------|
| Reactivity           | Human;Mouse;Rat                                     |
| Immunogen            | Recombinant Human MMP-2 protein                     |
| Host                 | Rabbit                                              |
| Isotype              | IgG                                                 |
| Clone                | C170083                                             |
| Purification         | Protein A                                           |
| Buffer               | 0.2 µm filtered solution in PBS                     |
| Applications         |                                                     |
| Recommended Dilution |                                                     |
| WB:                  | 1:1000-1:5000; ICC/IF: 1:100-1:200; FC: 1:200-1:500 |
| Data                 |                                                     |



Western blot detection of MMP2 in mouse lung, mouse heart cell lysates using MMP2 antibody (1:1000 diluted).

**Observed-MW: 72,64 kDa**

**Calculated-MW: 74 kDa**

| Preparation & Storage |                                                                 |
|-----------------------|-----------------------------------------------------------------|
| Storage               | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping              | Ice bag                                                         |

**Background**

This gene is a member of the matrix metalloproteinase (MMP) gene family, that are zinc-dependent enzymes capable of cleaving components of the extracellular matrix and molecules involved in signal transduction. The protein encoded by this gene is a gelatinase A, type IV collagenase, that contains three fibronectin type II repeats in its catalytic site that allow binding of denatured type IV and V collagen and elastin. Unlike most MMP family members, activation of this protein can occur on the cell membrane. This enzyme can be activated extracellularly by proteases, or, intracellularly by its S-glutathiolation with no requirement for proteolytical removal of the pro-domain. This protein is thought to be involved in multiple pathways including roles in the nervous system, endometrial menstrual breakdown, regulation of vascularization, and metastasis. Mutations in this gene have been associated with Winchester syndrome and Nodulosis-Arthropathy-Osteolysis (NAO) syndrome. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Oct 2014]

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.5