



A Reliable Research Partner in Life Science and Medicine

# **Human PⅢNP Antibody Pair SetSet**

Catalog No. E-KAB-0164 Applications ELISA

Synonyms P3NP, N-Propeptide Of Type III Procollagen, Procollagen III Amino Terminal Propeptide

## **Kit components & Storage**

Title	Specifications	Storage
Human PⅢNP Capture Antibody	1 vial, 100 μ g	Store at -20℃. Avoid freeze/thaw
		cycles.
Human PIIINP Detection Antibody	1 vial, 50 μL	Store at -20°C. Avoid freeze/thaw
(Biotin)		cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0164)	
		Human PIIINP Capture Antibody	Human PⅢNP Detection Antibody
			(Biotin)
Immunogen	Immunogen	Native Protein	Native Protein
Information	Swissprot	P02461	
Product details	Reactivity	Human	Human
	Host	Mouse	Mouse
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300; 1%
		glycerol, pH 7.4	protective protein; 50% glycerol; pH
			7.4
	Purify	Protein A	Protein A
	Specificity	Detects Human PIIINP in ELISAs.	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>



A Reliable Research Partner in Life Science and Medicine

## **Applications**

Human PIIINP Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Human PⅢNP Capture Antibody	
Capture			.10
			Arro
ELISA	1:1000-1:10000	Human PIIINP Detection Antibody	Optical Desity
	1.1000-1.10000	•	5 1.1
Detection		(Biotin)	
			0.01
			Human PIIINP concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

### **Background**

This gene encodes the pro-alpha1 chains of type III collagen, a fibrillar collagen that is found in extensible connective tissues such as skin, lung, uterus, intestine and the vascular system, frequently in association with type I collagen. Mutations in this gene are associated with Ehlers-Danlos syndrome types IV, and with aortic and arterial aneurysms. Two transcripts, resulting from the use of alternate polyadenylation signals, have been identified for this gene.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>