

A Reliable Research Partner in Life Science and Medicine

# **APC Anti-Human CD11a Antibody[HI111]**

Catalog Number: E-AB-F1316E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human Host Mouse

**Isotype** Mouse IgG1, κ

Clone No. HI111

Isotype Control APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]

Conjugation APC

**Conjugation Information** APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### **Data**



Human peripheral blood lymphocytes are stained with APC Anti-Human CD11a Antibody[HI111] (filled gray histogram) or APC Mouse IgG1, κ Isotype Control (empty black histogram).

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

#### **Antigen Information**

Alternate Names CD11 antigen-like family member A;CD11a;Integrin alpha-L;Itgal;LFA-1A;Ly-15;

Lymphocyte antigen 15

 Uniprot ID
 P20701

 Gene ID
 3683

## For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

#### **Background**

Elabscience®

CD11a is a 170-180 kD type I transmembrane glycoprotein also known as LFA-1a chain and integrin αL subunit. CD11a non-covalently associates with integrin β2 (CD1 8) to form LFA-1. It is expressed on all leukocytes, including B and T lymphocytes, monocytes, macrophages, neutrophils, basophils and eosinophils. It is absent on nonhematopoietic tissues and platelets. CD11a plays a central role in leukocyte cell-cell interactions and is important in lymphocyte costimulation. CD11a/CD18 binds to ICAM-1 (CD54), ICAM-2 (CD102), and ICAM-3 (CD50).

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Email:techsupport@elabscience.com Web:www.elabscience.com